

STUDIES ON THE EFFECT OF BLANCHING MEDIA ON YIELD AND NUTRITIONAL QUALITY OF CARROT JUICE

Zia-ur-Rehman and Wazir Hussain Shah*

Biotechnology & Food Research Centre, PCSIR Laboratories Complex, Lahore-54600, Pakistan

(Received 1 July 1996; accepted 7 January 1998)

Effect of various blanching processes on the retention of different nutrients in carrot juice products was investigated. Maximum brix value of the juice was 9.3 when carrots were blanched in 0.05 N acetic acid solution containing 0.2% CaCl₂. However, brix values of the juice obtained from unblanched and water blanched carrots were 8.1 and 8.5 respectively. Highest amount of carotene and vitamin C was also found in the juice which was obtained from carrots after blanching in acetic acid and 0.2% CaCl₂ solution. Other blanching processes showed adverse effects on the retention of carotene in carrot juice products compared with unblanched carrot juice.

Key words: Carrot, Blanching processes, Carrot juice, Nutritional value.

Introduction

Carrot (*Daucus carota* L.) is a rich source of carotene, soluble sugars and minerals (Bajaj *et al* 1980). Apart from these nutrients, carrot also contains sufficient amount of dietary fibre (Robertson *et al* 1979). Although carrot is consumed as vegetable in Pakistan but its juice is also used as soft drink due to its delicious taste and high nutritional value. Different types of beverages based on carrot juice are also being prepared in many areas of the world (Lombran and Dias 1985; Kim and Gerber 1988). In order to extract maximum juice from carrots, different blanching processes and extracting techniques have been tried during the past few years (Stephens *et al* 1976; Jan 1987; Sims *et al* 1993). It has been observed by many workers that blanching process improves the extractability of juice from carrots to some extent (Khan *et al* 1975; Bao and Chang 1994). However, very little information is available in the literature regarding the effect of blanching on the nutritional quality of carrot juice. This communication reports the effect of different blanching processes on the availability of various nutrients in carrot juice.

Materials and Methods

Juice extraction. Fresh mature carrots of red variety, free from blemishes, were washed thoroughly and peeled after removing their heads and tails with a sharp knife manually. The carrots were cut longitudinally into two halves and then blanched according to the method of Stephens *et al* (1976). 5 kg cleaned carrots were treated for 5 min in 12.5 l of boiling solution in each blanching material reported in Table 2,

thoroughly drained and then ground through a champion juicer (Plastaket Manufacturing Co, Lodi CA) equipped with a stainless steel screen.

Analysis. pH of the juice was determined using a glass electrode pH meter (PYE Unicam, England). Titratable acidity of the samples was estimated by titration against 0.1 NaOH solution using phenolphthalein as indicator (AOAC 1990). Brix of juice samples was measured with a Bausch and Lomb refractometer. Carotene was estimated using the method of Valaden and Mummery (1975) after extracting it in petroleum ether and acetone mixture. Vitamin C content in carrot juice was determined on spectrophotometer according to Bajaj and Gurdeep method (1981). Total soluble sugars of carrot pulp products were estimated according to Dinitrosalicylic acid method as described by Miller (1959).

Statistical analysis. All data were subjected to analysis of variance and standard deviation was calculated by the method of Steel and Torrie (1980).

Results and Discussion

Chemical composition of dried carrots is given in Table 1. The amount of proteins, lipids and minerals in dried carrots was 6.64, 2.98 and 7.69% respectively. Significant amount of total water soluble nutrients (49.8%) which mainly consisted of soluble sugars (30.27%) were also present in carrots. Apart from these nutrients, carrots also contained considerable amount of carotene and vitamin C contents.

Effect of blanching on the yield of carrot juice. Table 2 shows the effect of blanching on the yield of juice from carrots. Fresh juice yield from unblanched carrots was 64.0% higher

* Author for correspondence

than the yield from blanched carrots in almost all the cases except one. Carrots blanched in acetic acid or citric acid produced 63.0 and 61.5% comparatively higher amount respectively than that of the water blanched carrots (54.6%). Similarly carrots blanched in KCl or CaCl₂ solutions also yielded more juice (57.0% and 59.1% respectively) than water blanched carrots but yielded lesser juice compared to carrots blanched in organic acid solutions. It seems that texture of whole carrots became soft after blanching, making juice extraction through the stainless steel screen more difficult as already reported by Sims *et al* (1993). However, maximum juice yield was 68.0% after blanching the carrots in acetic acid solution containing 0.2% CaCl₂. It seems that solubility characteristics of pectic substances were considerably increased due to greater diffusion of calcium into the carrots in the presence of acetic acid which consequently increased the extractability of juice upto 6%. However, exact mechanism is still unknown. Juice and pulp collection processes caused 1.4-4.5% losses which were comparatively less than those reported by earlier workers (Bao and Chang 1994).

Table 1
Chemical composition* of dried carrot

Parameters		
1.	Protein %	664 (± 1.5)
2.	Lipids %	2.98 (± 1.0)
3.	Minerals (Ash) %	7.69 (± 2.3)
4.	Total soluble nutrients %	49.8 (± 1.9)
5.	Total soluble sugars %	30.2 (± 1.8)
6.	Reducing sugars %	9.1 (± 0.9)
7.	Non-reducing sugars %	21.1 (± 1.3)
8.	Carotene (mg 100g ⁻¹)	17.6 (± 1.6)
9.	Vitamin C (mg 100g ⁻¹)	45 (± 1.9)

* Average of three replicates along with standard deviation.

Effect of blanching on pH, acidity and brix of juice. pH of carrot juice extracted from unblanched carrots was 6.22, whereas it was 6.18 when extracted from water blanched carrots (Table 3). The pH value of the juice extracted from carrots blanched in KCl or CaCl₂ solutions remained almost unchanged. However, a significant decrease in pH of the juice was observed when carrots were blanched in acidic solutions. Titratable acidity of the juice extracted from blanched carrots was higher than the juice of unblanched carrots. The increase in titratable acidity and decrease in pH was probably due to heat decomposition of pectic substances into pectic acid and absorption of organic acids by carrots (Stephens *et al* 1971). The changes in pH and titratable acidity were found significant ($P < 0.05$).

Blanching process also affected the brix of the juice (Table 3). Brix of the juice obtained from unblanched and water blanched carrots was 8.1 and 8.5 respectively. The brix of juices obtained from carrots blanched in acetic acid with or without 0.2% CaCl₂ was 9.3 and 8.9 respectively; whereas the brix values of the juices extracted from those carrots blanched in citric acid with or without 0.2% CaCl₂ were also higher than unblanched carrot juice. The results were in agreement with the findings of Stephens *et al* (1971) who reported that the juice from acetic acid blanched carrots was slightly higher in brix than juice from carrots blanched in water. It seems that combined treatment of acetic acid or citric acid with CaCl₂ retarded tissue softening effect and cleaved glycosidic as well as ionic linkages of pectin molecules, resulting in greater water solubility of pectic substances, which may be responsible for increasing brix values of the juice.

Effect of blanching on carotene and vitamin C contents of juice. Juice from unblanched carrots had carotene contents (170 mg 100g⁻¹) higher than the juice extracted from blanched carrots (Table 3). However, the juice extracted from the

Table 2
Effect of blanching on the yield of carrot (%)* juice

Blanching treatment	Juice	Pulp	Loss
1. Unblanched	64.0 (0.3)	12.2 (0.2)	3.8(0.4)
2. Distilled water	54.6 (0.3)	43.0 (0.4)	2.4 (0.3)
3. 0.2% KCl solution	57.8 (0.6)	40.0 (0.5)	3.0 (0.1)
4. 0.2% CaCl ₂ solution	59.1 (0.4)	38.2 (0.3)	2.7 (0.2)
5. 0.05N Acetic acid solution	63.0 (0.7)	33.0 (0.5)	4.0 (0.3)
6. 0.5N Acetic acid + 0.2% CaCl ₂ solution	68.0 (0.9)	30.6 (0.6)	1.4 (0.3)
7. 0.05N Citric acid solution	61.5 (1.2)	34.0 (0.9)	4.5 (0.4)
8. 0.05N Citric acid + 0.2% CaCl ₂ solution	64.8(0.9)	32.0 (0.5)	3.2 (0.2)

* Average of three replicates along with standard deviation.

Table 3
Effect of blanching on pH, acidity, brix, carotene and vitamin C contents of carrot juice

Blanching treatment	pH	Titrateable acidity (%)	Brix	Carotene (mg 100g ⁻¹)	Vitamin C (mg 100g ⁻¹)
1. Unblanched	6.22(0.01)	0.08(0.02)	8.1(0.1)	170(1.2)	5.00(0.6)
2. Distilled water	6.18(0.01)	0.11(0.01)	8.5(0.1)	155(1.1)	6.66(0.9)
3. 0.2% KCl solution	6.20 (0.02)	0.09 (0.02)	8.3 (0.2)	150 (1.7)	6.90 (0.4)
4. 0.2% CaCl ₂ solution	6.20 (0.04)	0.10 (0.01)	8.2 (0.3)	153 (1.3)	6.90 (0.6)
5. 0.05N acetic acid solution	5.95 (0.02)	0.15 (0.03)	8.0 (0.2)	167 (1.0)	7.00(0.7)
6. 0.05N Acetic acid + 0.2% CaCl ₂ solution	5.85(0.01)	0.14(0.01)	9.3(0.3)	195(1.2)	8.44(0.9)
7. 0.05N Citric acid solution	5.90(0.02)	0.13(0.02)	8.7(0.2)	165(1.9)	7.00(0.7)
8. 0.05N Citric acid + 0.2% CaCl ₂ solution	5.90(0.02)	0.13(0.02)	9.1(0.4)	169(1.3)	7.96(0.9)

carrots blanched in acetic acid containing 0.2% CaCl₂ retained the highest amount of carotene (195 mg 100g⁻¹) among all other juices (P<0.05). This might be due to easy extractability of carotene from the tissues. It is also possible that carotene was being protected from destruction in the presence of acetic acid and CaCl₂ during high temperature blanching process. Other blanching processes reduced the carotene in the juice to some extent.

It is apparent from Table 3 that blanching processes showed significant effect on the extraction of vitamin C in carrot juice products. Fresh juice from unblanched carrots had the lowest vitamin C retention (5.0 mg 100g⁻¹). However, vitamin C contents in juice obtained from blanched carrots varied from 6.66 to 8.44 mg 100g⁻¹. Juice obtained from carrots blanched in acetic acid containing 0.2% CaCl₂ had the highest amount of vitamin C (8.44 mg 100g⁻¹). It could be the result of easy extraction of vitamin C from the carrot tissues.

Conclusion

It is apparent from these studies that different blanching processes affected the extractability of juice from carrots. About 6% more juice was extracted on blanching the carrots in 0.05N acetic acid solution containing 0.2% CaCl₂. Retention of carotene and vitamin C was also the highest in this blanching process. Brix value of unblanched carrot juice was 8.1, whereas 9.3 was the brix value of the juice obtained after blanching carrots in acetic acid solution containing 0.2% CaCl₂. Other blanching processes showed adverse effects on the extraction of juice and retention of carotene in carrot juice products.

References

- A O A C 1990 Official Methods of Analysis, Association of Official Analytical Chemists, 15th ed, Washington, D.C.
- Bajaj K L, Gurdeep K, Sukhija B S 1980 Chemical composition and some plant characteristic in relation to quality of some promising cultivars of carrots (*Daucus carota* L.). *Qual Plant Plant Foods Hum Nutr* **30** 97-107.
- Bajaj K L, Gurdeep K 1981 Spectrophotometric determination of L-ascorbic acid in vegetables and fruits. *Analyst* **106**, 1117-1120.
- Bao B, Chang K C 1994 Carrot juice color, carotenoids and non starchy polysaccharides as affected by processing conditions. *J Fd Sci* **59** 1155-1158.
- Jun H 1987 Bioavailability of carotene in carrot juice. *Dissertation Abstracts International B-47* 3300.
- Khan N A, Khalil J K, Khan I 1975 Changes in the carotene content of carrots during maturity, processing and storage under different temperature conditions. *Pak J Sci Res* **27** 112-114.
- Kim H, Gerber L H 1988 Influence of processing on quality of carrot juice. *Korean J Fd Sci Technol* **20** 683-686.
- Lombran J I, Dias J M 1985 Rheological and chemical changes in stored carrot juice. *Can Inst Food Sci Technol* **18** 213-219.
- Miller G L 1959 Use of dinitrosalicylic acid reagent for determination of reducing sugars. *Anal Chem* **31** 426-428.
- Roberts J A, Eastwood M A, Yeoman M M 1979 An investigation into the dietary fibre content of named varieties of carrots at different development ages. *J Sci Fd Agric* **30** 388-394.
- Sims C A, Balsban M O, Mathews R F 1993 Optimization of carrot juice and cloud stability. *J Fd Sci* **58** 1129-1131.

- Steel R G, Torrie J H 1980 *Principles and Procedures of Statistics*. McGraw Hill, London pp 345.
- Stephens T S, Saldana G, Brown H E, Griffiths F P 1971 Stabilization of carrot juice by dilute acid treatment. *J Fd Sci* **36** 36-38.

- Stephens T S, Saldana G, Lime B J 1976 Neutralized juice of acid treated carrots. *J Fd Sci* **41** 1245-1246.
- Valadon L P G, Mummery R S 1975 Carotenoids of floral plants and the spadix of *Arum maculatum*. *Z Pflanzen Physiol* **75** 88-94.