

Antibacterial Activity of Artemisia

Malika Ait Ouaddi*, Meryem Elgarini and Mohammed Sobh

Laboratory of Health, Environment and Biotechnology, Group of Physiopathology, Molecular Genetic and Biotechnology, Faculty of Sciences Ain Chock, University Hassan-II, Casablanca, Morocco

(received October 2, 2023; revised May 1, 2024; accepted June 3, 2024)

Abstract. Currently, bacterial resistance to antibiotics has become a real problem in the world, which has pushed scientific research towards the search for alternatives represented by secondary metabolites synthesized by medicinal and aromatic plants, especially essential oils. The present study is a bibliographical synthesis characterizing the chemical composition of essential oils as well as the antibacterial activity of five species belonging to the 'Artemisia' genus: *A. absinthium*, *A. mésoatlantica*, *A. annua*, *A. herba alba* et *A. campestris*, against gram positive and gram-negative bacteria most resistant to antibiotics generally used in medicine. The study carried out gave the following results: the majority of gram-positive bacteria such as *Staphylococcus* and *Streptococcus* and gram-negative bacteria such as extended-spectrum beta-lactamase enterobacteriaceae: *Escherichia coli*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* have become resistant or even multi-resistant to greater or lesser percentages with conventional antibiotics, even with third generation antibiotics. The essential oils of the plants studied had a very significant effect against all gram-positive bacteria: *Staphylococcus aureus*, *Listeria monocytogenes*, *Bacillus subtilis*, *Micrococcus luteus*, *Bacillus cereus* and *Streptococcus* as well as against gram negative bacteria: *Escherichia coli*, *Salmonella*, *Klebsiella*, *Pseudomonas aeruginosa* and *Proteus mirabilis*. These oils are generally characterized by their richness in powerful antibacterial substances, especially those belonging to oxygenated monoterpenes. Antibacterial activity is also related to stereochemistry, synergy effect and type of bacteria. Indeed, the essential oils of medicinal and aromatic plants are an excellent natural alternative to the antibiotics currently used in medicine.

Keywords: antibacterial activity, antibiotic, artemisia, essential oil, monoterpenes

Introduction

Public health suffers, especially in recent decades from several infectious diseases whose causative agents have become resistant or even multi-resistant to existing antibiotics WHO (1999). This has caused interested parties to start looking for alternatives to deliver antimicrobial molecules to overcome the problem of bacterial resistance to antibiotics. Medicinal plants, especially those used traditionally, are the main source of these antimicrobial molecules Bouharb *et al.* (2014). Morocco is characterized by its wealth of medicinal and aromatic plants, hence the economic importance of this sector in the country. Morocco exports the equivalent of 250 million dirhams in PAM to the United States and the European Union USAID (2005). Economically, essential oils bring in almost 165 million dirhams USAID (2005). Artemisia is a genus of medicinal and aromatic plants belonging to the Asteraceae family and which is characterized by a very significant chemical polymorphism. This chemo variety

can be observed from one stand to another or even between two plants of the same species. This influences the yield and chemical quality of the essential oil from these plants Abad *et al.* (2012). Species belonging to the genus Artemisia are generally located in the north in temperate regions, while their distribution in southern regions remains limited Oberprieler *et al.* (2009). The genus Artemisia includes about 500 species distributed between herbaceous plants and shrubs. These species can be divided into 5 or 6 groups or subgenera given the great polymorphism that exists within this genus. The 6 representative species of these different subgenera are: *Artemisia annua* and *Artemisia vulgaris* of the subgenus Artemisia, *Artemisia absinthium* of the subgenus Absinthium, *Artemisia tridentata* of the subgenus Tridentata, *Artemisia dracunculus* of the subgenus Dracunculus and *Artemisia chinensis* of the subgenus Pacifica Vallès *et al.* (2011). In Morocco, Artemisia is represented by 19 species, the most important of which are: *Artemisia herba alba*, *Artemisia chitachensis*, *Artemisia negrei*, *Artemisia ifranensis*, *Artemisia arborescens* and *Artemisia absinthium* Ibn Tatou and Fennane (2008).

*Author for correspondence;
E-mail: malikaouaddi4@gmail.com

Bacterial resistance to antibiotics. Antibiotic resistance has become a real problem in recent decades, especially with the overuse of antibiotics. These can no longer kill certain bacteria or slow down their growth. This resistance concerns both gram-positive and gram-negative bacteria.

Gram positive bacteria. The bacteriology laboratory of the Mohammed V military Hospital of Rabat in Morocco carried out a study between April 2012 and March 2016 on the resistance of gram-positive bacteria to antibiotics. *Staphylococcus aureus* was resistant to methicillin in 8% of cases. Methicillin resistant *Staphylococcus aureus* isolates were resistant to gentamicin with a percentage of 48%, Ciprofloxacin (37.5%), Tetracycline (50.6%) and Tobramycin (52.6%). No isolate was resistant to Glycopeptides. Resistance to Fusidic acid, Rifampicin, Fosfomycin was respectively 48%, 15%, 17%. For *Streptococcus*, all species remain highly sensitive to antibiotics, except *Streptococcus B*, which are 90% resistant to Tetracycline.

Gram negative bacteria. The majority of *Enterococcus* isolates were represented by *Escherichia* species. *Faecalis* at 90.5%, with resistance to tetracycline at 78.5%, ampicillin at 1.6% and rifampicin at 12.5%. *Escherichia faecium* was resistant to 80% erythromycin and 68% lincomycin. No isolate was resistant to Glycopeptides Lahlou *et al.* (2020). ESBL enterobacteriaceae represented mainly by *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter cloacae*, *Enterobacter aerogenes*, *Klebsiella oxytoca*, *Citrobacter koseri*, *Citrobacter braakii* and *Serratia marcescens* showed an antibiotic resistance rate of 60.14%: 89.13% to Norfloxacin, 87% to Ciprofloxacin, 50% to Gentamicin, 11% to Amikacin, 80.43% to trimethoprim-sulfamethoxazole combination and 2.17% to Imipenem. The strains studied were all sensitive to colistin. ESBL *Escherichia coli* had a resistance rate of 42% to gentamicin, 74.19% to the trimethoprim-sulfamethoxazole combination, 16.12% to Amikacin and 87.09% to ciprofloxacin and norfloxacin. The strains studied were sensitive to imipenem and colistin. The ESBL *Klebsiella pneumonia* resistance rate was 100% to norfloxacin, 93% to ciprofloxacin as well as to the trimethoprim-sulfamethoxazole combination, 64.28% to gentamicin, 7.14% to imipenem and all strains were sensitive to colistin and amikacin. The resistance rate of *Pseudomonas aeruginosa* is 100% to aztreonam, to ticarcillin-clavulanic acid combination, as well as to ticarcillin. The strains studied were all sensitive to colistin. Their

resistance to ceftazidime was 10% with a resistance rate of 66% to amikacin, gentamicin, ciprofloxacin, Imipenem, as well as to the combination Piperacillin-tazobactam Amhal (2017).

The antibacterial activity of Artemisia. *Artemisia absinthium*. *Artemisia absinthium* or wormwood is a perennial shrub plant that has great medical interests, it is native to Asia, middle east, Europe and north Africa Sharopov *et al.* (2012). It contains many bioactive phytochemicals such as: terpenoids (trans-thujones, γ -terpinene, 1.4 terpineol, myrcene, bornyl acetate, cadinene, camphene, trans-sabinyacetate, guaiazulene, chamazulene, camphor and linalool), lactones, organic acids, tannins, resins and phenols Omer *et al.* (2007). It also contains flavonoids, glycosides and phenolic acids Kordali *et al.* (2005). The essential oil of *Artemisia absinthium* is rich in myrcene, trans-thujones, cis-epoxyocimene, cis chrysanthenyl acetate and trans-sabinyl acetate which are the compounds generally most found in this species Dhen *et al.* (2014); Rezaeinodehi and Khangholi (2008). Absinthe is also known by absinthin, substances that characterize this species. The fresh plant is an important source of azulene El Saber Batiha *et al.* (2020); Bhat *et al.* (2019).

Ethanol extract of *Artemisia absinthium* inhibits *Staphylococcus aureus* ATCC29213 with a zone of inhibition between 10 and 15 mm in diameter. They have no remarkable activity against *Escherichia coli*, *Staphylococcus faecalis* and *Bacillus subtilis* var niger ATCC10 Dulger *et al.* (1999). *Artemisia absinthium* had a significant effect against gram-positive pathogenic bacteria Fiamegos *et al.* (2011). The methanolic extracts of *Artemisia absinthium* inhibit *Bacillus subtilis* ATCC6633, *Salmonella typhimurium* RSSK95091, *Bacillus cereus* 6230, thermophilic *Salmonella* 6453, *Providencia alcalifaciens* 3215 and *Pseudomonas putida* 1617. Moslemi *et al.* (2012) that the hydroalcoholic extract has an antibacterial effect against *Staphylococcus aureus*. The antibacterial activity noted may be due to the synergistic effect between the majority compounds (camphor, cymene...etc) and the minority compounds (α -pinene, β - pinene etc.) contained in the essential oil. This activity is strongly linked to the bioactive compounds which are the hydrocarbon monoterpenes, which they are (α -pinene, β - pinene, camphor, caryophyllene and p-cymene. These molecules make the biological membrane of bacteria permeable Moslemi *et al.* (2012). The antibacterial action of essential oil extracted from *Artemisia absinthium* was more accentuated than

gentamicin against *Staphylococcus aureus* (the resistant and sensitive strains), *Salmonella typhimurium*, *Escherichia coli* ATCC8739, *Proteus vulgaris*, *Klebsiella pneumoniae* 10031 and *Pseudomonas aeruginosa* 9027. This activity is related to the presence of α -phellandrene and chamazulene which are the important components of essential oil Assili *et al.* (2015); Rezaeinodehi and Khangholi (2008). Mihajiliv and Krstevetal recorded that the minimum inhibitory concentration of *Artemisia absinthium* essential oil was less than 0.08 mg/mL for *Staphylococcus aureus* 25923 and *Pseudomonas aeruginosa* 9027 extracted from wounds and for *Pseudomonas mirabilis* extracted from human stool and 2.43 mg/mL for *Klebsiella oxytoca* isolated from stool and 0.14 μ L/mL for *Listeria monocytogenes*. These actions are related to the suppression of protein, polysaccharide, DNA and RNA biosynthesis in bacterial cells. (Table 1) Mihajilov-Krstev *et al.* (2014); Belay *et al.* (2011).

Artemisia mesatlantica. The study of the essential oil from *Artemisia mesatlantica* collected in the region of Ifrane in Morocco revealed the presence of terpene esters with terpinyl acetate, terpene alcohols (α -terpineol), monoterpenes (camphene, limonene, myrcene, tri-cyclene), oxides with 1.8 cineole, sesquiterpene ketones (piperidone) and sesquiterpenes. The essential oil of *Artemisia mesatlantica* is characterized by the dominance of β -thujone followed by camphene and camphor. In addition to the majority compounds, there are other minority compounds such as 1.8 cineole, D-germacrene, 1-epicubenol, α -thujone and davanone. The classification of these molecules according to their functional chemical groups shows the predominance of oxygenated monoterpenes Bencheqroun *et al.* (2012).

For antibacterial activity, the essential oil of this plant has an antibacterial effect against *Escherichia coli* and *Staphylococcus aureus* with a dilution of 1/2000 v/v, *Bacillus subtilis* with a dilution of 1/3000 v/v and *Micrococcus luteus* with a dilution of 1/1000 v/v (Table 2) Bencheqroun *et al.* (2012).

Artemisia annua. The essential oil of *Artemisia annua*, collected in the southern slopes of the middle Atlas in Morocco, is rich in bioactive molecules such as:

Table 2. Antibacterial activity of the essential oil of *Artemisia mesatlantica*

Bacteria	<i>E. coli</i>	<i>B. subtilis</i>	<i>M. luteus</i>	<i>S. aureus</i>
Dilution(v/v)	1/2000	1/3000	1/1000	1/2000

oxygenated monoterpenes (1.8 cineole, artemisia ketone, artemisia camphor alcohol, borneol), hydrocarbon sesquiterpenes (copaene, γ -cadinene, caryophyllene, β -farnesene, germacrene D and β -selinene), oxygenated sesquiterpenes (caryophyllene oxide).

Chebbak *et al.* (2023) showed that *Artemisia annua* studied has a significant antibacterial effect against gram-positive bacteria but especially against gram negative bacteria especially *Escherichia coli* with a zone of inhibition close to 75.67 mm and a minimum inhibitory concentration of 5.34 μ L/mL plus or minus 0.50 μ L/mL followed by *Bacillus subtilis* with an inhibition zone of 37 mm and minimum inhibitory concentration of 5.64 μ L/mL plus or minus 0.97 μ L/mL then *Staphylococcus aureus* with a zone of inhibition close to 23.33 mm and a minimum inhibitory concentration equal to 4.98 μ L/mL plus or minus 1.82 μ L/mL, then *Klebsiella pneumoniae* with a zone of inhibition close to 22.33 mm and a minimum inhibitory concentration 5.44 μ L/mL followed by *Salmonella sp* with a zone of inhibition close to 16 mm and a minimum inhibitory concentration of 1.09 mg/mL and finally *Bacillus cereus* which is the least sensitive bacterium with an inhibition zone of 11 mm and a minimum inhibitory concentration of 5.44 mg/mL. (Table 3) Chebbak *et al.* (2023).

Artemisia herba alba. The essential oil of *Artemisia herba-alba* of Moroccan origin has been studied from a chemical point of view and antibacterial power. This essential oil is composed of: oxygenated monoterpenes (56.4%) followed by oxygenated sesquiterpenes (2.5%). The other constituents represent 38.7%. the components found are: cis-thujones, trans-thujones, vanillyl alcohol, davanone, cis-threo-davanafuran, camphor, isobornyl *n*-butyrate, cis-chrysanthenyl acetate, trans-arbusculone, jasmolactone, *n*-dodecanol, artedouglasia oxide C and cyclosativene. *Artemisia herba-alba* essential oil had

Table 1. Antibacterial activity of the essential oil of *Artemisia absinthium* Mihajiliv -Krstev *et al.* (2014)

Bacteria	<i>S. aureus</i> 25923	<i>P. aeruginosa</i> 9027	<i>P. mirabilis</i>	<i>K. oxytoca</i>	<i>L. monocytogenes</i>
MIC (mg/mL)	0.08	0.08	0.08	2.43	0.14

Table 3. Antibacterial activity of the essential oil of *Artemisia annua*

Bacteria	Zone of inhibition (mm)	MIC(μ L/mL)
<i>E. coli</i>	75.67	5.34
<i>B. subtilis</i>	37	5.64
<i>S. aureus</i>	23.33	4.98
<i>K. pneumoniae</i>	22.33	5.44
<i>Salmonella sp.</i>	16	1.09 mg/mL
<i>B. cereus</i>	11	5.44 g/mL

antibacterial activity against 12 bacterial species from the most sensitive to the least sensitive: *Bacillus clausii* 2226, *Brochotrix thermosphacta* D274, *Staphylococcus sp* GB1, *Streptococcus salivarius*, *Salmonella typhimurium*, *Carnobacterium maltaromaticum* 9P, *Pseudomonas fragi* 6P2, *Staphylococcus sp* ES1, *Serratia proteamaculans* 20P, *Brochotrix thermosphacta* 7R1. A very weak effect on *Escherichia coli* 32 and on *Staphylococcus aureus* Amor *et al.* (2019).

Artemisia herba-alba from other countries and whose main constituents are: cis and trans-thujones have shown the same results Majdouli and Elazzouzi (2015); Sbayou and Ababou (2014).

Artemisia herba-alba is characterized by significant chemical polymorphism. Thus, several chemotypes have been studied such as the eucalyptol chemotype and the caryophyllene acetate chemotype found in Jordan Aldosary *et al.* (2018); Qnais *et al.* (2016). Camphor has been noted as the main component of *artemisia herba alba* from Algeria and Tunisia Bertella and Benlahcen (2018); Younsi *et al.* (2017).

***Artemisia campestris*.** The volatile compounds contained in the essential oil of *Artemisia campestris* belong to four groups: oxygenated monoterpenes (49.5%), hydrocarbon monoterpenes (42.2%), oxygenated sesquiterpenes (2.9%) and hydrocarbon sesquiterpenes (2.8 %). The other constituents represent only 0.2%. the isolated components are diverse. Among them are: α -pinene (18.4%), camphene (7.7%), limonene (5.2%), camphor (9.2%), borneol (5.2%), α -terpineol (3.6%), α -terpinyl acetate (18.8%), bornyl acetate (2.7%), α -thujene (0,3%), trans- β -ocimene (2%) Belhattab *et al.* (2011). The constituents of essential oil of *artemisia campestris* differ according to the chemotype or variety. However, generally, the constituents in common and which characterize this species are: β -pinene (24,2_27,9%), p-cimene (17,4_22,3%) et α -pinene (4,1_11%) Gucker and Coreyl (2007).

The antibacterial activity of *artemisia campestris* essential oil is tested against *Escherichia coli* ATCC 25922, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* 27853, *Salmonella typhimurium*, *Staphylococcus aureus* ATCC 43300 and *Staphylococcus aureus*. The most potent antibacterial effect was observed against *Pseudomonas aeruginosa* ATCC 27853 with a zone of inhibition of 23 mm and against *Escherichia coli* with a zone of inhibition of 20 mm Djidel and Khenouf (2014).

Factors influencing antibacterial activity. In general, the presence of a significant concentration of oxygenated monoterpenes like thujones, camphor, camphene and 1.8-cineole in the essential oil increases its antibacterial activity. This may explain the pronounced effect of essential oil like that of *artemisia mesatlantica* against *Staphylococcus aureus* and *Bacillus subtilis* because *Staphylococcus aureus* has been shown to be most affected by ketone monoterpenes like thujones Oussalah, *et al.* (2007); Dorman and Deans (2000).

Other studies have shown that the majority of essential oil tested for their antibacterial effects had more pronounced activity against gram positive bacteria compared to gram negative bacteria, this is due to the structure of the outer membrane of gram-negative bacteria which is hydrophilic which prevents the entry of volatile water insoluble compounds Wan (1998).

Antibacterial activity is also influenced by stereochemistry. Studies by Dorman and Deans in 2000 that β -isomer compounds are more active against bacteria than α -isomers Dorman and Deans (2000). Synergy is a factor that also influences antibacterial activity. Indeed, the molecular interaction of the functional groups of different compounds of the essential oil with the wall of the bacteria leads to strong lesions. The cooperation therefore between the different majority and minority compounds of essential oils reinforces their antibacterial power Felice *et al.* (2004).

Conclusion

The present work has made it possible to study some species belonging to the genus *artemisia* with the aim of drawing conclusions that may be useful in the medical field. Indeed, *artemisia* is characterized, in general, by a powerful antibacterial power, whether against gram positive or gram-negative bacteria. This allows it to become a good alternative to current antibiotics in the medical and pharmaceutical field and specific formulas

that bring together specific compounds that can have a significant effect depending on the type of bacteria.

References

- Abad, M.J., Bedoya, L.M., Apaza, L., Bermejo, P. 2012. The *Artemisia* L. Genus: a review of bioactive essential oils. *Molecules*, **17**: 2542-2566.
- Aldosary, N.H., Omar, D., Awang, R.M., Adam, N.A. 2018. Chemical profiling and insecticidal activity of artemisia Herba-Alba essential oil against papaya mealybug *Paracoccus marginatus* (Hemiptera: Pseudococcidae). *Research Journal of Applied Sciences, Engineering and Technology*, **15**: 261-269.
- Al-Zubairi, A.S., Abdul, A.B., Abdel wahab, S.I., Peng, C.Y., Mohan, S., Elhassan, M.M. 2011. Eleusine indica possesses antioxidant, antibacterial and cytotoxic properties. *Evidence Based Complement Alternate Medicine*, **2011**: 965370.
- Amhal, F.Z. 2017. Profil épidémiologique Actuel Des Bactéries Multirésistantes Expérience de L'Hôpital Militaire Avicenne de Marrakech. *Ph.D. Thesis*, 142 pp., Faculty of Medicine and Pharmacy, Marrakech, Morocco.
- Amor, G., Caputo, L., La Stora, A., De Feo, V., Mauriello, G., Fechtali, T. 2019. Chemical composition and antimicrobial activity of *Artemisia herba-alba* and *Origanum majorana* essential oils from Morocco. *Molecules*, **24**: 4021.
- Asili, J., Emami, S.A., Einolghozati, R., Noghab, Z.S., Bazzaz, B.S.F., Sahebkar, A. 2015. Chemical composition and *in vitro* efficacy of essential oil of seven *Artemisia* species against ESBL producing *Escherichia coli* multidrug-resistant. *Journal of Essential Oil Bearing Plants*, **18**: 124-145.
- Belay, G., Tariku, Y., Kebede, T., Hymete, A., Mekonnen, Y. 2011. Ethnopharmacological investigations of essential oils isolated from five Ethiopian medicinal plants against eleven pathogenic bacterial strains. *Phytopharmacology*, **1**: 133-143.
- Belhattab, R., Boudjouref, M., Barroso, J.G., Pedro, L.P., Figueirido, A.C. 2011. Essential oil composition from *Artemisia campestris* grow in Algeria. *Advances in Environmental Biology*, **5**: 429-432.
- Bencheqroun, H.K., Ghanmi, M., Satrani, B., Aafi, A. 2012. Activité antimicrobienne des huiles essentielles d'*Artemisia mesatlantica*, plante endémique du Maroc. *Bulletin de la Société Royale des Sciences de Liège*, **81**: 4-21.
- Bertella, A., Benlahcen, K. 2018. *Artemisia herba-alba* Asso: essential oil antibacterial activity and acute toxicity. *Industrial Crops and Product*, **116**: 137-143.
- Bhat, R.R., Rehman, M.U., Shabir, A., Mir, M.U.R., Ahmad, A., Khan, R., Masoodi, M.H., Madkhali, H., Ganaie, M.A. 2019. Chemical composition and biological uses of *Artemisia absinthium* (wormwood). In: *Plant and Human Health*, vol. **3**, pp. 37-63, Springer, Berlin/Heidelberg, Germany.
- Boucharb, H., El Badaoui, K., Zair, T., El Amri, J., Chakir, S., Alaoui, T. 2014. Sélection de quelques plantes médicinales du Zerhoun (Maroc central) pour l'activité antibactérienne contre *Pseudomonas aeruginosa*. *Journal of Applied Bioscience*, **78**: 6685-6693.
- Benziane, O.Z., Eluoussaoui, A., Chalka, M., Lafraxo, S., Bin Jardan, Y.A., Bourhia, M., Guemmouh, R., Nafidi, H.A., Chebbak, K. 2023. Antimicrobial and antioxidant properties of chemically analyzed essential oil of *Artemisia annua* L. (Asteraceae) native to mediterranean area. *Life*, **13**: 807.
- Dahanukar, S.A., Kulkarni, R.A., Rege, N.N. 2000. Pharmacology of medicinal plants and natural products. *Indian Journal of Pharmacology*, **32**: 81-118.
- Dhen, N., Majdoub, O., Souguir, S., Tayeb, W., Laarif, A., Chaieb, I. 2014. Chemical composition and fumigant toxicity of *Artemisia absinthium* essential oil against *Rhyzopertha dominica* and *Spodoptera littoralis*, Tunis. *Journal of Plant Protect*, **9**: 57-61.
- Djidjel, S., Khennouf, S. 2014. Radical scavenging reducing power, lipid peroxidation inhibition and chelating properties of extracts from *Artemisia campestris* L. aerial parts. *Annual Research and Review in Biology*, **4**: 1691-170.
- Dorman, H.J.D., Deans, S.G. 2000. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, **88**: 308-316.
- Dülger, B., Ceylan, M., Alitsaous, M., U-gurlu, E. 1999. Antimicrobial Activity of *Artemisia absinthium* L. *Turkish Journal of Biology*, **23**: 377-384.
- Ekwenye, U.N., Elegalam, N.N. 2005. Antibacterial activity of ginger (*Zingiber officinale* Roscoe) and garlic (*Allium sativum* L.) extracts on *Escherichia coli* and *Salmonella typhi*. *International Journal of Molecular Medicine and Advance Sciences*, **1**: 411-417.
- El-Mleeh, A., Hetta H.F., Al-Rejaie, S., Alghamdi, S., Zahoor, M., Magdy, B.A., Murata, T., Zaragoza-Bastida, A., Rivero-Perez, N., El saber Batiha, G., Olatunde, A. 2020. Bioactive compounds, Pharmacological actions and pharmacokinetics of wormwood (*A. absinthium*). *Antibiotics*, **2020**: 9-353.
- Felice, S., Francesco, N., Nelly, A.A., Maurizio, B.,

- Werner, H. 2005. Composition and antimicrobial activity of the essential oil of *Achillea falcata* L. (Asteraceae). *Flavour and Fragrance Journal*, **20**: 291-294.
- Fiamegos, Y.C., Kastritis, P.L., Exarchou, V., Han, H., Bonvin, A.M., Vervoort, J., Lewis, K., Hamblin, M.R., Tegos, G.P. 2011. Antimicrobial and efflux pump inhibitory activity of caffeoylquinic acids from *Artemisia absinthium* against gram-positive pathogenic bacteria. *PLOS ONE*, **6**: 18127.
- Gucker, Corey, L. 2007. *Artemisia campestris*. US Department of agriculture, Forest Service, Rocky Mountain Research Station, Retrieved 2008 from <http://www.Fs.Fed.US/database/feis>.
- Ibn Tattou, M., Fennane, M. 2008. Flore vasculaire du Maroc : inventaire et chorologie. *Travaux Institut Scientifique Rabat série Botanique*, **2**: 39-398.
- Kordali, S., Cakir, A., Mavi, A., Kilic, H., Yildirim, A. 2005. Screening of chemical composition and antifungal and antioxidant activities of the essential oils from three Turkish *Artemisia* species. *Journal of Agriculture and Food Chemistry*, **53**: 1408-1416.
- Lahlou, Y.B., Comlan, A.G., Maleb, A., Lemnouer, A., Elouennass, M. 2020. Épidémiologie et profil de résistance des bactéries à gram positif à l'hôpital militaire d'instruction Mohammed v de Rabat, Maroc. *Journal Marocain des Sciences Médicales*, **22**: 20-24.
- Majdoui, K., Soro, N.K., Elazzouzi, H., Benmaachou, J., Drioiche, A., Zaïr, T. 2015. Chemical composition and antibacterial activity of *Artemisia Herba Alba Huguetii* essential oil from south of Morocco. *International Journal of Current Research*, **7**: 14397-14404.
- Mihajilov-Krstev, T., Jovanoviæ, B., Joviæ, J., Iliæ, B., Miladinoviæ, D., Matejiæ, J., Rajkoviæ, J., Ðorðeviæ, L., Cvetkoviæ, V., Zlatkoviæ, B. 2014. Antimicrobial, antioxidative and insect repellent effects of *Artemisia absinthium* essential oil. *Planta Medica*, **80**: 1698-1705.
- Moslemi, H.R., Hoseinzadeh, H., Badouei, M.A., Kafshdouzan, K., Fard, R.M.N. 2012. Antimicrobial activity of *Artemisia absinthium* against surgical wounds infected by *Staphylococcus aureus* in a rat model. *Indian Journal of Microbiology*, **52**: 601-604.
- Oberprieler, C., Himmelreich, S., Källersjö, M., Vallès, J., Watson, L.E., Vogt, R. 2009. Anthemideae. *Systematics, Evolution and Biogeography of the Compositae*, **38**: 632-666.
- Omer, B., Krebs, S., Omer, H., Noor, T. 2007. Steroid-sparing effect of wormwood (*Artemisia absinthium*) in Crohn's disease: a double-blind placebo-controlled study. *Phytomedicine*, **14**: 87-95.
- Oussalah, M., Caillet, S., Saucier, L., Lacroix, M. 2007. Inhibitory effects of selected plant essential oils on the growth of four pathogenic bacteria: *E. coli* O157: H7, *Salmonella Thyphimurium*, *Staphylococcus aureus* and *Listeria monocytogenes*. *Food Control*, **18**: 414-420.
- Qnais, E.Y., Alatshan, A.Z., Bseiso, Y.G. 2016. Chemical composition, antinociceptive and anti-inflammatory effects of *Artemisia herba-alba* essential oil. *Journal of Food, Agriculture and Environment*, **14**: 20-27.
- Rezaeinodehi, A., Khangholi, S. 2008. Chemical composition of the essential oil of *Artemisia absinthium* growing wild in Iran. *Pakistan Journal of Biological Science*, **11**: 946-949.
- Sbayou, H., Ababou, B. 2014. Chemical composition and antibacterial activity of *Artemisia herba-alba* and *Mentha pulegium* essential oils. *Journal of Life Science*, **8**: 35-41.
- Sharopov, F., Sulaymonova, V.A., Setzer, W.N. 2012. Composition of the essential oil of *Artemisia absinthium*. *Records of Natural Products*, **6**: 127-134.
- Tan, R.X., Zheng, W.F., Tang, H.Q. 1998. Biologically active substances from the genus *Artemisia*. *Planta Medica*, **64**: 295-302.
- USAID. 2005. Chemonics international, inc. filière des plantes aromatiques médicinales. U.S. Agency for International Development (USAID)/Morocco.
- Vallès, J., Garcia, S., Hidalgo, O., Martín, J., Pellicer, J., Sanz, M., Garnatje, T.M. 2009. Biology, genome evolution, biotechnological issues and research including applied perspectives in *Artemisia* (Asteraceae). *Advances in Botanical Research*, **60**: 350-384.
- Wan, J., Wilcock, A., Coventry, M.J. 1998. The effect of essential oils of basil of the growth *Aeromonas hydrophila* and *Pseudomonas fluorescens*. *Journal of Applied Microbiology*, **84**: 152-158.
- Guidelines Subcommittee. 1999. 1999 World Health Organization-International Society of Hypertension Guidelines for the Management of Hypertension. *Journal of Hypertension*, **17**: 151-183.
- Younsi, F., Mehdi, S., Aissi, O., Rahali, N., Jaouadi, R., Boussand, M., Messaoud, C. 2017. Essential oil variability in natural populations of *Artemisia campestris* (L.) and *Artemisia herba-alba* (Asso) and incidence on antiacetylcholinesterase and antioxidant activities. *Chemical Biodiversity*, **14**: 1700017.